

# ***MIT Joint Program on the Science and Policy of Global Change***



## **Unintended Environmental Consequences of a Global Biofuels Program**

*Jerry M. Melillo, Angelo C. Gurgel, David W. Kicklighter, John M. Reilly,  
Timothy W. Cronin, Benjamin S. Felzer, Sergey Paltsev, C. Adam Schlosser,  
Andrei P. Sokolov, and X. Wang*

**Report No. 168  
January 2009**

The MIT Joint Program on the Science and Policy of Global Change is an organization for research, independent policy analysis, and public education in global environmental change. It seeks to provide leadership in understanding scientific, economic, and ecological aspects of this difficult issue, and combining them into policy assessments that serve the needs of ongoing national and international discussions. To this end, the Program brings together an interdisciplinary group from two established research centers at MIT: the Center for Global Change Science (CGCS) and the Center for Energy and Environmental Policy Research (CEEPR). These two centers bridge many key areas of the needed intellectual work, and additional essential areas are covered by other MIT departments, by collaboration with the Ecosystems Center of the Marine Biology Laboratory (MBL) at Woods Hole, and by short- and long-term visitors to the Program. The Program involves sponsorship and active participation by industry, government, and non-profit organizations.

To inform processes of policy development and implementation, climate change research needs to focus on improving the prediction of those variables that are most relevant to economic, social, and environmental effects. In turn, the greenhouse gas and atmospheric aerosol assumptions underlying climate analysis need to be related to the economic, technological, and political forces that drive emissions, and to the results of international agreements and mitigation. Further, assessments of possible societal and ecosystem impacts, and analysis of mitigation strategies, need to be based on realistic evaluation of the uncertainties of climate science.

This report is one of a series intended to communicate research results and improve public understanding of climate issues, thereby contributing to informed debate about the climate issue, the uncertainties, and the economic and social implications of policy alternatives. Titles in the Report Series to date are listed on the inside back cover.


Henry D. Jacoby and Ronald G. Prinn,  
*Program Co-Directors*

For more information, please contact the Joint Program Office

Postal Address: Joint Program on the Science and Policy of Global Change  
77 Massachusetts Avenue  
MIT E19-411  
Cambridge MA 02139-4307 (USA)

Location: 400 Main Street, Cambridge  
Building E19, Room 411  
Massachusetts Institute of Technology

Access: Phone: (617) 253-7492  
Fax: (617) 253-9845  
E-mail: [globalchange@mit.edu](mailto:globalchange@mit.edu)  
Web site: <http://globalchange.mit.edu/>

 Printed on recycled paper

# Unintended Environmental Consequences of a Global Biofuels Program

Jerry M. Melillo<sup>\*ψ</sup>, Angelo C. Gurgel<sup>#</sup>, David W. Kicklighter<sup>\*</sup>, John M. Reilly<sup>#</sup>, Timothy W. Cronin<sup>\*</sup>, Benjamin S. Felzer<sup>\*</sup>, Sergey Paltsev<sup>#</sup>, C. Adam Schlosser<sup>#</sup>, Andrei P. Sokolov<sup>#</sup> and X. Wang<sup>§</sup>

## Abstract

*Biofuels are being promoted as an important part of the global energy mix to meet the climate change challenge. The environmental costs of biofuels produced with current technologies at small scales have been studied, but little research has been done on the consequences of an aggressive global biofuels program with advanced technologies using cellulosic feedstocks. Here, with simulation modeling, we explore two scenarios for cellulosic biofuels production and find that both could contribute substantially to future global-scale energy needs, but with significant unintended environmental consequences. As the land supply is squeezed to make way for vast areas of biofuels crops, the global landscape is defined by either the clearing of large swathes of natural forest, or the intensification of agricultural operations worldwide. The greenhouse gas implications of land-use conversion differ substantially between the two scenarios, but in both, numerous biodiversity hotspots suffer from serious habitat loss. Cellulosic biofuels may yet serve as a crucial wedge in the solution to the climate change problem, but must be deployed with caution so as not to jeopardize biodiversity, compromise ecosystems services, or undermine climate policy.*

## Contents

1. INTRODUCTION.....	1
2. METHODS .....	2
3. RESULTS AND DISCUSSION .....	4
3.1 Changes in the Global Landscape – How Land is Used .....	4
3.2 Carbon Balance .....	5
3.3 Impacts on Biodiversity.....	7
3.4 A Larger Human Footprint on the Land .....	9
4. CONCLUSIONS.....	10
5. REFERENCES.....	11
6. APPENDIX.....	15
A.1 The Terrestrial Ecosystem Model (TEM).....	15
A.2 MIT Emissions Predictions and Policy Analysis (EPPA) Model.....	16
A.3 Downscaling EPPA Estimates of Land-use Change.....	19
A.4 Mapping EPPA Land Shares to TEM Cohorts .....	21
A.5. Development of the Climate Scenario .....	22

## 1. INTRODUCTION

At the end of the 20<sup>th</sup> century, the world’s commercial energy consumption was about 400 exa-joules (EJ) per year, with fossil fuels contributing about 85% and all others (nuclear, biofuels, hydro, wind, solar) contributing only 15%. Typical projections of the world economy

---

<sup>\*</sup> The Ecosystems Center, Marine Biological Library, Woods Hole, MA

<sup>#</sup> Joint Program on the Science and Policy of Global Change, MIT, Cambridge MA

<sup>§</sup> Parsons Brinckerhoff, Inc., Boston, MA

<sup>ψ</sup> Corresponding author: Jerry M. Melillo (Email: jmelillo@mbl.edu)

imply energy demands in 2050 of 550-1000 EJ per year, depending on resource availability, and the price, scope and effect on energy demand of policies to limit greenhouse gas (GHG) emissions and air pollutants (Clarke *et al.*, 2007). To limit GHG emissions, we will need a variety of low-carbon energy sources operating at very large scales; for example, sources supplying 55-100 EJ/year would meet only about 10% of the estimated demand. Biofuels are being promoted as an important part of the global energy mix in the coming decades to meet the climate change challenge (Pacala and Socolow, 2004; Farrell *et al.*, 2006).

Recently, there has also been considerable emphasis on the social and environmental costs of current biofuels technologies (Tilman *et al.*, 2006; Fargione *et al.*, 2008; Scharlemann and Laurance, 2008; Searchinger *et al.*, 2008; The Royal Society, 2008). Increased production of biofuel crops has the potential to compete with food production for arable land. In addition, increased biofuels production could require conversion of natural lands with resulting carbon emissions, threats to biodiversity, and possible likely increased use of fertilizers and pesticides. At the same time, a growing population will create increasing demand for food, while changes in the climate, CO<sub>2</sub> and tropospheric ozone will affect land requirements and the location of production activities. To date, most of the analyses of environmental impacts of biofuels have been done as local or sub-regional case studies without explicit consideration of concurrent environmental changes and growing demands for food. In addition, they make very simple assumptions about which land types are converted for biofuels production (Searchinger *et al.*, 2008).

Here, we use a computable general equilibrium (CGE) model of the world economy, the MIT Emissions Predictions and Policy Analysis model (EPPA, Paltsev *et al.*, 2005; Gurgel *et al.*, 2007), coupled with a process-based terrestrial biogeochemistry model, the Terrestrial Ecosystem Model (TEM, Melillo *et al.*, 1993; Felzer *et al.*, 2004; Sokolov *et al.*, 2008), to generate global land-use scenarios and to explore some of the environmental consequences of an aggressive global cellulosic biofuels program over the first half of the 21<sup>st</sup> century. The biofuels scenarios we focus on are linked to a global climate policy to control greenhouse gas emissions from industrial and fossil fuel sources that would, absent feedbacks from land-use change, stabilize the atmosphere's CO<sub>2</sub> concentration at 550 ppmv (Paltsev *et al.*, 2008). The climate policy makes the use of fossil fuels more expensive and speeds up the introduction of biofuels,

and ultimately increases the size of the biofuel industry, with additional effects on land use, land prices, and food and forestry production and prices.

## 2. METHODS

The amount of land used for biofuels production in a region depends not only on the ability of local environmental conditions to support crop productivity at a sufficient level, but also depends on competitive demands on the land to provide adequate food and fiber for the local population or to provide products for global trade. To examine the ability of terrestrial ecosystems to supply biofuels to meet a growing global demand for energy along with growing demands for food and fiber, we have developed an approach that links TEM to EPPA to generate world-wide land-use future scenarios at a spatial resolution of  $0.5^\circ$  latitude x  $0.5^\circ$  longitude that vary with climate change.

In our approach, greenhouse gas emissions, as projected by EPPA, drive a coupled atmospheric and climate module within the MIT Integrated Global System Model (IGSM, Prinn *et al.*, 1999; Sokolov *et al.*, 2005) to simulate the future climate that then drives TEM. What results of these model linkages is a set of projected changes in crop, pasture, and forest productivity as simulated in TEM due to changing climate, levels of CO<sub>2</sub> and tropospheric ozone. These projected changes in productivity are then fed back to the EPPA model to change yields in the agricultural sectors (Reilly *et al.*, 2007a). Changes in yields, together with changing demand for these products, as driven by population and income growth, lead to reallocations of land among uses, and conversions of land among land types. The regionally aggregated land-use types used by EPPA are downscaled to the  $\frac{1}{2}$  by  $\frac{1}{2}$  degree grid level based on a statistical approach for use in TEM (Wang, 2008). This linked modeling process generates scenarios that capture first-order interactions among land use, climate, and the economy. The pattern of land use is affected by a number of factors including population and economic growth, changing climate, and atmospheric concentrations of CO<sub>2</sub> and tropospheric ozone as they concurrently affect both overall productivity and the regional pattern of production. In addition, climate policy and energy demand affect land use as they drive demand for biofuels. The TEM is then used to evaluate the magnitude of GHG emissions from the land associated with these land-use changes projected by EPPA such as the conversion of forests to areas that produce biofuels. Additional details about our approach of coupling TEM to EPPA are provided in the Appendix.

For this study, we develop two scenarios that have the same economic growth and meet the same limit on industrial and fossil fuel GHG emissions. One scenario makes all land available for biofuels crops or other managed uses as long as the economic return on the land exceeds the cost of conversion and improvement. The other scenario limits access to unmanaged (e.g., tropical forests), with the limits based on the recent history of regional land conversion rates. This approach results in slower rates of deforestation than would be predicted by cost estimates alone (Gurgel *et al.*, 2007).

We refer to the first scenario as the “deforestation scenario” because it involves large-scale deforestation in support of biofuels production, either directly or indirectly. The direct link between deforestation and biofuels is when forests are cleared to establish biofuels crops (Fargione *et al.*, 2008). The indirect link is when biofuels production moves on to croplands or pastures, and causes new forest clearing to relocate agriculture (Searchinger *et al.*, 2008). We call the second scenario the “intensification scenario” because one possible result of limited access to new land is that existing managed lands will be used more intensively, with increased inputs of capital, labor and materials such as fertilizers. For each scenario, the initial land cover distribution is based on the land cover distribution for the year 2000, which has been derived by reorganizing the gridded land-use transitions data sets of Hurtt *et al.* (2006) for use by TEM and EPPA.


### **3. RESULTS AND DISCUSSION**

Energy from cellulosic biofuels plays an important part in the global primary energy supply in 2050 in both scenarios – 141 EJ yr<sup>-1</sup> in the deforestation scenario and 128 EJ yr<sup>-1</sup> in the intensification scenario, with these levels of energy supply large enough to meet at least 10% of the projected global energy requirement in 2050. Our simulations with the two scenarios explore how the production of these large amounts of energy affects several features of the global environment. The features we consider include land area devoted to producing cellulosic biofuels, carbon storage on land and biodiversity, especially in the sub-tropics and tropics.

#### **3.1 Changes in the Global Landscape – How Land is Used**

At the beginning of the 21<sup>st</sup> century about 31.5% of the total land area of 133 million km<sup>2</sup> was in agriculture; 12.1% (16.1 million km<sup>2</sup>) in crops and 19.4% (25.8 million km<sup>2</sup>) in pasture (Hurtt *et al.*, 2006), with no land devoted to cellulosic biofuels. Land used to produce the feedstocks for

the current generation of biofuels (e.g., ethanol from maize (corn) and sugarcane) is included in the crop sector. In the deforestation scenario, we estimate that by 2050, the land area in cellulosic biofuels will grow to 14.8 million km<sup>2</sup>, which is 11.1% of the earth's total land area (**Figure 1a**). At the same time, we project that the area of croplands will grow to about 20.0 million km<sup>2</sup> and the area in pasture will shrink slightly to 24.5 million km<sup>2</sup>. The growth of croplands by 3.9 million km<sup>2</sup> over the first half of the 21<sup>st</sup> century in the deforestation scenario is in response to increased food demands globally.



**Figure 1.** Distribution of cellulosic biofuels in 2050 simulated for two scenarios: deforestation (a) and intensification (b). Data expressed as the percentage of each ½ by ½ degree grid cell devoted to biofuels production.

By 2050 in the intensification scenario, we estimate that the land area in cellulosic biofuels will grow to 13.9 million km<sup>2</sup> (**Figure 1b**). We also project that by mid-century croplands area will grow by about 2 million km<sup>2</sup> to almost 18 million km<sup>2</sup> (**Table 1**), and pasture areas will shrink by almost 8 million km<sup>2</sup> to just under 18 million km<sup>2</sup>. Despite the loss of pastures, the intensification of land use allows nearly as much production of food as in the deforestation scenario.

**Table 1.** Areas of land-use changes associated with crops, pastures and biofuels over the first half of the 21<sup>st</sup> century. The estimated land surface of the earth is estimated to be 133.02 million km<sup>2</sup>.

	<b>Absolute land areas in different land uses (million km<sup>2</sup>)</b>		
	<b>Current</b>	<b>Deforestation Scenario</b>	<b>Intensification Scenario</b>
Year	2000	2050	2050
Total Land Area of the Earth	133.02	133.02	133.02
Total Area Co-opted for Human Use	41.96	59.29	49.84
Biofuels Area	0.00	14.79	13.91
Crop Area	16.12	20.01	17.99
Pasture Area	25.84	24.49	17.94

### 3.2 Carbon Balance

Energy production from biofuels together with agriculture expansion will result in a large loss in carbon from land ecosystems as natural vegetation such as forests and savannas is cleared to grow cellulosic feedstocks, crops and pastures (**Table 2**). Such a reduction in terrestrial carbon storage associated with the expansion of biofuels and other land uses has been referred to as a “carbon debt” (Fargione *et al.*, 2008; Searchinger *et al.*, 2008). Through time, this debt can be canceled if biofuels production and use has net carbon (and other greenhouse gas) emissions that are less than the total emissions of the fossil fuels they displace. The overall “carbon balance” is thus initially negative, due to the carbon debt incurred by land conversion, but slowly rises as the annual credits from the biofuels repay (and eventually outweigh) the one-time debt.

**Table 2.** Effects of intensive management (food crops, pastures and biofuels) on the global carbon cycle simulated for two scenarios, deforestation and intensification, during the first half of the 21<sup>st</sup> century.

<b>Land Cover</b>	<b>Deforestation Scenario</b>		<b>Intensification Scenario</b>	
	<b>ΔCarbon (Pg C)</b>	<b>Co-opted NPP (Pg C yr<sup>-1</sup>)</b>	<b>ΔCarbon (Pg C)</b>	<b>Co-opted NPP (Pg C yr<sup>-1</sup>)</b>
Time Period	2000-2050	2040-2049	2000-2050	2040-2049
Biofuel Crops	-21.38	8.21	+4.34	7.18
Food Crops	-53.53	10.49	-18.73	9.63
Pasture	-28.23	10.29	-19.25	7.55
Total Agriculture	-103.14	28.99	-33.64	24.36
Percent Co-opted	--	50.09	--	42.09


The total carbon debt associated with biofuels is the sum of the direct and indirect carbon debts – direct debt consisting only of losses from the lands used for biofuels, and indirect debt




consisting of losses due to displacement of food-crops and pastures. The expansion of food-crop agriculture and pastures as projected in our simulation is a complex response to several factors including a growing demand for food as the human population increases, competition for fertile lands to grow biofuels feedstocks, and reduced food crop yields per unit area in some regions as a consequence of climate change and increased tropospheric ozone pollution (Reilly *et al.*, 2007a). Due to these complex interactions, the indirect carbon debt associated with biofuels is extremely difficult to quantify, but we can consider carbon losses from all food crops and pastures to be an upper bound.

In the deforestation scenario, we project a direct carbon debt of 21 Pg C by 2050, with much of this carbon coming from areas once covered by tropical forests in Brazil and in Southeast Asia (**Figure 2a**). Indirect carbon debt could be as large as 82 Pg C, giving a total carbon debt of 21-103 Pg C (Table 2). This carbon debt is equal to 8-37% of the cumulative fossil fuel emissions for the period 2000-2050 in the climate policy we impose here to limit these emissions. Even considering the best-case, where total carbon debt is 21 Pg C, we estimate that the carbon debt associated with biofuels establishment in the deforestation scenario will last until the middle of the 21<sup>st</sup> century; that is, no net greenhouse gas reductions will be realized from biofuels until about 2045 (**Figure 3a**). Moving towards the worst-case, where total carbon debt is 103 Pg C, it becomes clear that these large emissions from land-use change would substantially undermine the efforts to stabilize climate.

The intensification scenario differs dramatically. Energy production from biofuels results in a direct carbon credit of 4 Pg C by 2050 (Table 1). The small carbon gains in many of the areas devoted to bioenergy production (**Figure 2b**) are mostly in soils in response to nitrogen fertilization, which stimulates plant growth and carbon inputs to soil. However, the indirect carbon debt is still potentially as large as 38 Pg C, giving a total carbon debt of -4 – 34 Pg C over the first half of the 21<sup>st</sup> century. While the upper bound on the total carbon debt is still substantial, it is much less than the 103 Pg C in the deforestation scenario (Table 1). This upper-bound debt of 34 Pg C in the intensification scenario is nearly repaid by the middle of the 21<sup>st</sup> century (**Figure 3b**).



**Figure 2.** Effects of cellulosic biofuels on terrestrial carbon storage over the first half of the 21<sup>st</sup> century as simulated for the deforestation scenario (a) and the intensification scenario (b). Negative numbers indicate a release of carbon from land ecosystems and positive numbers indicate an accumulation of carbon by land ecosystems.




**Figure 3.** Carbon balance associated with all land use change and that directly associated with biofuels over the period 2000-2050 as simulated by the deforestation (a) and intensification (b) scenarios. Carbon balance is calculated as the savings associated with the substitution of biofuels for fossil fuels minus the carbon released from the land used for biofuels (direct effects) along with food crops and pastures (indirect effects). Negative numbers indicate a net loss of carbon and positive numbers indicate a net gain.

### 3.3 Impacts on Biodiversity

One of the great challenges with cellulosic biofuels is developing ways to produce large quantities of plant feedstocks in a region without destroying its biodiversity. Evaluation of potential impacts should include both the direct effects of converting land to biofuels production and the indirect effects of the displacement of food crops and pastures to new areas as described


earlier. In both the deforestation and intensification scenarios, we project that by the middle of the 21<sup>st</sup> century, many regions will substantially increase the fraction of land they devote to meeting the combined demands for food and biofuels at the expense of natural ecosystems including a number of biodiversity “hotspots” in the sub-tropics and tropics. To qualify as a hotspot, a region must meet two strict criteria: it must contain at least 1,500 species of vascular plants (> 0.5 percent of the world’s total) as endemics, and it has to have lost at least 70 percent of its original habitat (Conservation International, 2008). Biodiversity hotspots currently at risk include the Mesoamerican forests, the cerrado of Brazil, the Guinean forests of West Africa, Madagascar, the Indo-Burma region of tropical Asia, and the forests of the Philippines, Malaysia and Indonesia in Southeast Asia (**Figure 4**).



**Figure 4.** Natural areas in many biodiversity hot spots have already been converted to crop and pasture agriculture and limited remaining areas would face more threats from biofuels expansion. Data shown is for circa 2000 and expressed as the percentage of each ½ by ½ degree grid cell devoted to the crop and pasture agriculture. Several biodiversity hotspots are circled – Mesoamerican forests (1); the cerrado of Brazil (2); the Guinean forests of West Africa (3); Madagascar (4); and the forests of Southeast Asia (5).

We project that the region devoting the largest area to biofuels production during 2050 will be sub-Saharan Africa – 5.2 million km<sup>2</sup> in the deforestation scenario and 5.7 million km<sup>2</sup> in the intensification scenario. In addition to this land requirement, the area devoted to food crops is projected to expand by 0.2 million km<sup>2</sup> by 2050 in the deforestation scenario, whereas it is projected to decrease by 0.4 million km<sup>2</sup> in the intensification scenario. In both scenarios, the

area of grazing lands is projected to decrease during the first part of the 21<sup>st</sup> century; by 0.5 million km<sup>2</sup> in the deforestation scenario and by 3.4 million km<sup>2</sup> in the intensification scenario. Because these changes in the area of agricultural lands are not enough to compensate for the area required for biofuels production, large tracts of natural forests, woodlands and grasslands will be converted to either food or cellulosic biofuels production. In the deforestation scenario (**Figure 5a**), we project the loss of over 3.1 million km<sup>2</sup> of the natural forest area, which means that about 59% of the natural forest in place at the start of the 21st century will be cleared by 2050. We also project a loss of 1.2 million km<sup>2</sup> of natural woodlands (a reduction of 63% compared to 2000). By contrast, in the intensification scenario (**Figure 5b**) we project that a smaller, but still substantial loss of 2.0 million km<sup>2</sup> of forest area by 2050; a reduction of 38% of the natural forest area in 2000. In addition, we estimate a loss of 0.7 million km<sup>2</sup> of natural woodlands (a reduction of 38%).



**Figure 5.** Loss of natural areas due to their conversion to crop and pasture agriculture and cellulosic biofuels between 2000 and 2050 as simulated by the deforestation (a) and intensification (b) scenarios. Data expressed as the percentage of each ½ by ½ degree grid cell devoted to agricultural and cellulosic biofuels production.

Latin America is the region with the second largest area devoted to cellulosic biofuels production, with 4.1 million km<sup>2</sup> in the deforestation scenario and 3.7 million km<sup>2</sup> in the intensification scenario by 2050. Also, in the deforestation scenario, the area devoted to grazing is projected to expand by 1.0 million km<sup>2</sup> between 2000 and 2050. These new requirements for managed lands are largely met through the clearing of species-rich forests and savannas (cerrado in Brazil). By 2050 in the deforestation scenario, we project the clearing of 5.2 million km<sup>2</sup> of natural forests (a reduction of 65%) and 0.6 million km<sup>2</sup> of natural woodlands (a reduction of 71%). Over the same period in the intensification scenario, we estimate the clearing of 1.6

million km<sup>2</sup> of natural forests (a reduction of 20%) and 0.2 million km<sup>2</sup> from savannas (a reduction of 20%). The reduced pressure on natural forests and savannas in the intensification scenario is due, in part, to the conversion of 0.4 million km<sup>2</sup> of managed forests producing woods products to areas producing biofuels.

Habitat destruction is a pervasive threat affecting biodiversity hotspots and is already causing extinctions in many areas. In both scenarios in our analysis, we project the loss of large areas of forest and savanna habitats due to the direct and indirect effects of implementing a large-scale biofuels program, although the areas lost are smaller in the intensification scenario. These losses have the potential to put thousands of endemic plant and animal species at risk across the globe, especially in the sub-tropical and tropical regions.

### **3.4 A Larger Human Footprint on the Land**

The fraction of terrestrial net primary production (NPP, the amount of new plant material produced each year) that human activities have appropriated for our purposes is an important index of the scale of human intervention in the biosphere (Vitousek *et al.* 1986, 1997). With our deforestation scenario, the amount of NPP directly co-opted by humans to grow crops, graze animals and produce biomass for biofuels rises in both the deforestation and intensification scenarios over the first five decades of the 21<sup>st</sup> century, but more in the former. At the start of the century, we estimate that about 32% of the terrestrial NPP is co-opted for agriculture – crops and pastures – which is in the range of 24 to 37% estimated by Haberl *et al.*, (2007). By mid-century we project this will increase to about 50% with the deforestation scenario and to about 42% with the intensification scenario with biofuels added to the sum (Table 1). Most of the increase in co-opted NPP by 2050 in each of the scenarios can be attributed to the production of biomass for cellulosic biofuels.

The increases in co-opted NPP coupled with the loss of biodiversity have the potential to diminish the capacity of terrestrial ecosystems to deliver many of the support services that humans rely on, such as the cleansing of air and water. We currently do not understand the relationships between ecosystem structure and function well enough to predict when such disturbances in a region will move it beyond a critical threshold for delivering one or more essential ecosystem service (Carpenter, 2003; Walker and Meyers, 2004; Millennium Ecosystem Assessment, 2005).

#### **4. CONCLUSIONS**

In our analyses, we have not accounted for all of the unintended consequences of intensification that are likely to occur and these additional effects are likely to differ between the two scenarios we have examined. Greater use of agricultural chemicals associated with intensification, such as fertilizers and pesticides, can lead to a variety of environmental problems. For example, additions of nitrogen fertilizer can result in the emission of nitrous oxide, a potent greenhouse gas, and the pollution of surface and ground water (Galloway *et al.*, 2003). Confined livestock production can lead to methane emissions from manure handling and disposal (Denman *et al.*, 2007). In addition, over grazing can lead to damage of riparian areas and cause soil erosion. How these problems are addressed in different regions of the world will eventually determine the environmental consequences of agriculture over the next half century. However, the addition of significant biofuels industry would create further pressures on the environment either through intensified use of existing land, or with more extensive use of land that would then require substantial conversion of natural lands.

Europe and the US have mandated significant use of biofuels in part because they are seen as reducing greenhouse gas emissions. Unfortunately, such technological-based policies often go awry because they fail to account for unintended environmental consequences. Existing and proposed emissions trading systems are, in principle, a superior approach for controlling greenhouse gases, but also fail to fully protect or provide incentive to increase carbon stocks in vegetation and soils. These poorly designed policies put carbon stocks in vegetation and soils at risk and in doing so potentially undermine the goal of stabilizing the atmospheric concentration of carbon dioxide at the desired target.

Even though we see the potential for considerable intensification of production on land, and therefore a less than one-for-one conversion of land to meet biofuels demands, the risks of converting land in biodiversity hotspots are substantial. With the loss of biodiversity comes a cascade of environmental consequences including the loss of critical ecosystem services (Millennium Ecosystem Assessment, 2005). It is clear that we must think holistically and proceed cautiously as we develop policies to use plant-based biofuels to combat global warming.

#### **5. REFERENCES**

Babiker, M., J. M. Reilly, M. Mayer, R. S. Eckaus, I. Sue Wing and R. Hyman, 2001: The MIT Emissions Prediction and Policy Analysis (EPPA) Model: Revisions, Sensitivities, and

- Comparison of Results. MIT JPSPGC, *Report 71*,  
[http://web.mit.edu/globalchange/www/MITJPSPGC\\_Rpt71.pdf](http://web.mit.edu/globalchange/www/MITJPSPGC_Rpt71.pdf)).
- Bonan, G. B., K. W. Oleson, M. Vertenstein, S. Lewis, X. Zeng, Y. Dai, R. E. Dickinson and Z-L Yang, 2002: The land surface climatology of the Community Land Model coupled to the NCAR Community Climate Model. *J. Climate*, **15**: 3123-3149.
- Brohan, P., J. J. Kennedy, I. Harris, S. F. B. Tett and P. D. Jones, 2006: Uncertainty estimates in regional and global observed temperature changes: a new dataset from 1850. *J. Geophysical Research*, **111**, D12106 (doi:10.1029/2005JD006548).
- Brovkin, V., M. Claussen, E. Driesschaert, T. Fichefet, D. Kicklighter, M. F. Loutre, H. D. Matthews, N. Ramankutty, M. Schaeffer and A. Sokolov, 2006: Biogeophysical effects of historical land cover changes simulated by six earth system models of intermediate complexity. *Climate Dynamics*, **26(6)**: 587-600 (doi:10.1007/s00382-005-0092-6).
- Clein, J. S., B. L. Kwiatkowski, A. D. McGuire, J. E. Hobbie, E. B. Rastetter, J. M. Melillo and D. W. Kicklighter, 2000: Modeling carbon responses of moist tundra ecosystems to historical and projected climate: A comparison of fine- and coarse-scale ecosystem models for identification of process-based uncertainties. *Global Change Biology*, **6(Suppl. 1)**: 127-140.
- Carpenter, S. R., 2003: *Regime Shifts in Lake Ecosystems: Pattern and Variation. Excellence in Ecology 15*, O. Kinne (ed.), International Ecology Institute: Germany.
- Clarke, L. E., J. A. Edmonds, H. D. Jacoby, H. M. Pitcher, J. M. Reilly and R. G. Richels, 2007: *Scenarios of Greenhouse Gas Emissions and Atmospheric Concentrations* (Sub-report 2.1A of Synthesis and Assessment Product 2.1 by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research, Dept. of Energy, Office of Biological & Environmental Research: Washington, DC, USA.
- Conservation International, 2008:  
<http://www.biodiversityhotspots.org/xp/Hotspots/pages/map.aspx>.
- Cramer, W. P., and R. Leemans, 2001: Global 30-Year Mean Monthly Climatology, 1930-1960, Version 2.1 (Cramer and Leemans). Data set. Available on-line [<http://www.daac.ornl.gov>] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A.
- Demographia, 2007: World Urban Areas: All Identified World Urbanized Areas of 500,000+ and Others—Population and Density (<http://www.demographia.com/db-worldua.pdf>).
- Denman, K. L., G. Brasseur, A. Chidthaisong, P. Ciais, P. M. Cox, R. E. Dickinson, D. Hauglustaine, C. Heinze, E. Holland, D. Jacob, U. Lohmann, S. Ramachandran, P. L. Da Silva Dias, S. C. Wofsy and X. Zhang, 2007: Couplings Between Changes in the Climate System and Biogeochemistry. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor and H. L. Miller (eds.), Cambridge University Press: Cambridge, United Kingdom & New York, NY, USA.
- Dimaranan, B., and R. McDougall, 2002: *Global Trade, Assistance and Production: The GTAP5 Data Base*. Center for Global Trade Analysis, Purdue University, West Lafayette, Indiana.
- Fargione, J., J. Hill, D. Tilman, S. Polasky and P. Hawthorne, 2008: Land clearing and the biofuel carbon debt. *Science*, **319**: 1235-1238 (doi:10.1126/science.1152747).
- Farrell, A. E., R. J. Plevin, B. T. Turner, A. D. Jones, M. O'Hare and D. M. Kammen, 2006: Ethanol can contribute to energy and environmental goals. *Science*, **311**: 506-508.



- Felzer, B., D. Kicklighter, J. Melillo, C. Wang, Q. Zhuang and R. Prinn, 2004: Effects of ozone on net primary production and carbon sequestration in the conterminous United States using a biogeochemistry model. *Tellus*, **56B**: 230-248.
- Felzer, B., J. Reilly, J. Melillo, D. Kicklighter, M. Sarofim, C. Wang, R. Prinn and Q. Zhuang, 2005: Future effects of ozone on carbon sequestration and climate change policy using a global biogeochemical model. *Climatic Change*, **73**: 345-373 (doi: 10.1007/s10584-005-6776-4).
- Felzer, B. S., T. Cronin, J. M. Reilly, J. M. Melillo and X. Wang, 2007: Impacts of ozone on trees and crops. *Comptes Rendus Geoscience*, **339**: 784-798.
- Galloway, J. N., J. D. Aber, J. W. Erisman, S. P. Seitzinger, R.W. Howarth, E. B. Cowling and B. J. Cosby, 2003: The nitrogen cascade. *BioScience*, **53(4)**: 341-356.
- Gurgel, A., J. Reilly and S. Paltsev, 2007: Potential land use implications of a global biofuels industry. *J. Agric. Food Industrial Org.*, **5(2)**: 1-34.
- Haberl, H., K. H. Erb, F. Krausmann, V. Gaube, A. Bondeau, C. Plutzer, S. Gingrich, W. Lucht and M. Fischer-Kowalski, 2007: Quantifying and mapping the human appropriation of net primary production in earth's terrestrial ecosystems. *Proc. Natl. Acad. Sci. USA*, **104**: 12942–12947 (doi: 10.1073/pnas.0704243104).
- Hertel, T., 1997. *Global Trade Analysis: Modeling and Applications*. Cambridge: Cambridge University Press.
- Hulme, M., T. J. Osborn and T. C. Johns, 1998: Precipitation sensitivity to global warming: Comparison of observations with HadCM2 simulations. *Geophys. Res. Letts.*, **25**: 3379-3382.
- Hurt, G. C., S. Frolking, M. G. Fearon, B. Moore, E. Shevliakova, S. Malyshev, S. W. Pacala and R. A. Houghton, 2006: The underpinnings of land-use history: three centuries of global gridded land-use transitions, wood harvest activity, and resulting secondary lands. *Global Change Biol.*, **12**: 1208-1229 (doi: 10.1111/j.1365-2486.2006.01150.x).
- Jacoby, H., R. Eckaus, A. D. Ellermann, R. Prinn, D. Reiner and Z. Yang, 1997: CO<sub>2</sub> emissions limits: Economic adjustments and the distribution of burdens. *The Energy Journal*, **18(3)**: 31-58; MIT JPSPGC, *Reprint 1997-1*.  
([http://globalchange.mit.edu/files/document/MITJPSPGC\\_Reprint97-1.pdf](http://globalchange.mit.edu/files/document/MITJPSPGC_Reprint97-1.pdf)).
- Lee, H., T. W. Hertel, B. Sohngen and N. Ramankutty, 2005: Towards an Integrated Land Use Data Base for Assessing the Potential for Greenhouse Gas Mitigation. GTAP Technical Paper 25, Center for Global Trade Analysis, Purdue University, West Lafayette, Indiana.
- Matthews, E., and I. Fung, 1987: Methane emission from natural wetlands: global distribution, area, and environmental characteristics of sources. *Global Biogeochemical Cycles*, **1(1)**: 61-86.
- McGuire, A. D., J. M. Melillo, D. W. Kicklighter, Y. Pan, X. Xiao, J. Helfrich, B. Moore III, C. J. Vorosmarty and A. L. Schloss, 1997: Equilibrium responses of global net primary production and carbon storage to doubled atmospheric carbon dioxide: Sensitivity to changes in vegetation nitrogen concentration. *Global Biogeochemical Cycles*, **11**: 173-189.
- McGuire, A. D., J. S. Clein, J. M. Melillo, D. W. Kicklighter, R. A. Meier, C. J. Vorosmarty and M. C. Serreze, 2000a: Modeling carbon responses of tundra ecosystems to historical and projected climate: Sensitivity of pan-arctic carbon storage to temporal and spatial variation in climate. *Global Change Biology*, **6(Suppl. 1)**: 141-159.



- McGuire, A. D., J. M. Melillo, J. T. Randerson, W. J. Parton, M. Heimann, R. A. Meier, J. S. Clein, D. W. Kicklighter and W. Sauf, 2000b: Modeling the effects of snowpack on heterotrophic respiration across northern temperate and high latitude regions: Comparison with measurements of atmospheric carbon dioxide in high latitudes. *Biogeochemistry*, **48**: 91-114.
- McGuire, A. D., S. Sitch, J. S. Clein, R. Dargaville, G. Esser, J. Foley, M. Heimann, F. Joos, J. Kaplan, D. W. Kicklighter, R. A. Meier, J. M. Melillo, B. Moore III, I. C. Prentice, N. Ramankutty, T. Reichenau, A. Schloss, H. Tian, L. J. Williams and U. Wittenberg, 2001: Carbon balance of the terrestrial biosphere in the twentieth century: analyses of CO<sub>2</sub>, climate and land-use effects with four process-based ecosystem models. *Global Biogeochemical Cycles*, **15**: 183-206.
- Melillo, J. M., A. D. McGuire, D. W. Kicklighter, B. Moore, C. J. Vorosmarty and A. L. Schloss, 1993: Global climate change and terrestrial net primary production. *Nature*, **363**: 234-240.
- Millennium Ecosystem Assessment, 2005: *Ecosystems and Human Well-being: Synthesis*. Island Press: Washington, DC.
- Millennium Ecosystem Assessment, 2005: [www.millenniumassessment.org/documents/document.300.aspx.pdf](http://www.millenniumassessment.org/documents/document.300.aspx.pdf).
- Pacala, S., and R. Socolow, 2004: Stabilization wedges: solving the climate problem for the next 50 years with current technologies. *Science*, **305**: 968-972.
- Paltsev, S., J. Reilly, H. Jacoby, A. Gurgel, G. Metcalf, A. Sokolov and J. Holak, 2008: Assessment of US Cap-and-Trade Proposals. *Climate Policy*, **8**(4): 395-420; see also MIT JPSPGC, *Report 146*, April, 95 p. ([http://web.mit.edu/globalchange/www/MITJPSPGC\\_Rpt146.pdf](http://web.mit.edu/globalchange/www/MITJPSPGC_Rpt146.pdf)).
- Paltsev, S., J. Reilly, H. Jacoby, R. Eckaus, J. McFarland and M. Babiker, 2005: *The MIT Emissions Prediction and Policy Analysis (EPPA) Model: Version 4* MIT JPSPGC, *Report 125*, August, 72 p. ([http://web.mit.edu/globalchange/www/MITJPSPGC\\_Rpt125.pdf](http://web.mit.edu/globalchange/www/MITJPSPGC_Rpt125.pdf)).
- Prinn, R., H. Jacoby, A. Sokolov, C. Wang, X. Xiao, Z. Yang, R. Eckhaus, P. Stone, D. Ellerman, J. Melillo, J. Fitzmaurice, D. Kicklighter, G. Holian and Y. Liu, 1999: Integrated global system model for climate policy assessment: Feedbacks and sensitivity studies. *Climatic Change*, **41**: 469-546.
- Reilly, J., S. Paltsev, B. Felzer, X. Wang, D. Kicklighter, J. Melillo, R. Prinn, M. Sarofim, A. Sokolov and C. Wang, 2007a: Global economic effects of changes in crops, pasture and forests due to changing climate, carbon dioxide and ozone. *Energy Policy*, **35**: 5370-5383.
- Reilly, J., B. Felzer, D. Kicklighter, J. Melillo, H. Tan and M. Asadoorian, 2007b. The Prospects for Biological Carbon Sinks in Greenhouse Gas Emissions Trading Systems. In: *Greenhouse Gas Sinks*, Reay *et al.* (eds.), CABI Publishing: Wallingford, UK, Chapter 8, pp. 115-142.
- Scharlemann, J.P.W. and W.F. Laurance, 2008: How green are biofuels? *Science*, **319**: 43-44, doi:10.1126/science.1153103.
- Schlosser, C. A., D. Kicklighter and A. P. Sokolov, 2007: A Land Model System for Integrated Global Change Assessments. MIT JPSPGC, *Report 147*, May, 58p. ([http://web.mit.edu/globalchange/www/MITJPSPGC\\_Rpt147.pdf](http://web.mit.edu/globalchange/www/MITJPSPGC_Rpt147.pdf)).
- Searchinger, T., R. Heimlich, R.A. Houghton, F. Dong, A. Elobeid, J. Fabiosa, S. Tokgoz, D. Hayes and T.-H. Yu, 2008: Use of U. S. croplands for biofuels increases greenhouse gases through emissions from land-use change. *Science*, **319**: 1238-1240, doi: 10.1126/science.1151861.

- Sohngen, B. 2007: *Global Timber Market and Forestry Data Project*: <http://aede.osu.edu/people/sohngen.1/forests/GTM/index.htm>.
- Sohngen, B., R. Mendelsohn, and R. Sedjo, 1999: Forest Management, Conservation, and Global Timber Markets. *American Journal of Agricultural Economics*, **81**: 1-13.
- Sokolov, A. P., D. W. Kicklighter, J. M. Melillo, B. S. Felzer, C. A. Schlosser and T. W. Cronin, 2008: Consequences of considering carbon-nitrogen interactions on the feedbacks between climate and the terrestrial carbon cycle. *Journal of Climate*, **21**: 3776-3796 (doi: 10.1175/2008JCLI2038.1).
- The Royal Society, 2008: *Sustainable Biofuels: Prospects and Challenges*. Science Policy Section, The Royal Society: London (<http://royalsociety.org/document.asp?id=7366>).
- Tian, H., J. M. Melillo, D. W. Kicklighter, A. D. McGuire, J. V. K. Helfrich III, B. Moore III and C. J. Vörösmarty, 1998: Effect of interannual climate variability on carbon storage in Amazonian ecosystems. *Nature*, **396**: 664-667.
- Tian, H., J. M. Melillo, D. W. Kicklighter, A. D. McGuire and J. Helfrich, 1999: The sensitivity of terrestrial carbon storage to historical climate variability and atmospheric CO<sub>2</sub> in the United States. *Tellus*, **51B**: 414-452.
- Tian, H., J. M. Melillo, D. W. Kicklighter, A. D. McGuire, J. Helfrich III, B. Moore III and C. J. Vörösmarty, 2000: Climatic and biotic controls on annual carbon storage in Amazonian ecosystems. *Global Ecology and Biogeography*, **9**: 315-336.
- Tian, H., J. M. Melillo, D. W. Kicklighter, S. Pan, J. Liu, A. D. McGuire and B. Moore III, 2003: Regional carbon dynamics in monsoon Asia and its implications for the global carbon cycle. *Global and Planetary Change*, **37**: 201-217 (doi:10.1016/S0921-8181(02)00205-9).
- Tilman, D., J. Hill and C. Lehman, 2006: Carbon-negative biofuels from low-input high-diversity grassland biomass. *Science*, **314**: 1598-1600 (doi: 10.1126/science.1133306).
- Vitousek, P. M., P. R. Ehrlich, A. H. Ehrlich and P. A. Matson, 1986: Human appropriation of the products of photosynthesis. *BioScience*, **36**: 368-373.
- Vitousek, P., H. A. Mooney, J. Lubchenco and J. M. Melillo, 1997: Human domination of the Earth's ecosystems. *Science*, **277**: 494-499.
- Xiao, X., D. W. Kicklighter, J. M. Melillo, A. D. McGuire, P. H. Stone, A. P. Sokolov, 1997: Linking a global terrestrial biogeochemical model and a 2-dimensional climate model: implications for the carbon budget. *Tellus*, **49B**: 18-37; MIT JPSPGC Reprint 1997-3.
- Xiao, X., J. M. Melillo, D. W. Kicklighter, A. D. McGuire, R. G. Prinn, C. Wang, P. H. Stone, A. Sokolov, 1998: Transient climate change and net ecosystem production of the terrestrial biosphere. *Global Biogeochemical Cycles*, **12**: 345-360.
- You, L., and S. Wood, 2006: An entropy approach to spatial disaggregation of agricultural production. *Agricultural Systems*, **90**: 329-347.
- Walker, B., and J. Meyers, 2004: Thresholds in ecological and social-ecological systems: A developing database. *Ecology and Society*, **9(2)**: 3 (<http://www.ecologyandsociety.org/vol9/iss2/art3/>).
- Wang, X., 2008: *Impacts of Greenhouse Gas Mitigation Policies on Agricultural Land*, Doctoral Thesis, Massachusetts Institute of Technology.
- Webster, M., C. Forest, J. Reilly, M. Babiker, D. Kicklighter, M. Mayer, R. Prinn, M. Sarofim, A. Sokolov, P. Stone and C. Wang, 2003: Uncertainty analysis of climate change and policy response. *Climatic Change*, **61(3)**: 295-320.

## 6. APPENDIX

Below, we first briefly describe TEM and EPPA along with recent modifications that enabled linkage between these two models. We then describe how land-use change estimates from EPPA are downscaled and organized for use in TEM. Finally, we describe the development of the future climate scenario and two future land-use change scenarios that allow us to explore alternative pathways for achieving a climate policy goal and the potential environmental consequences of those pathways.

### A.1 The Terrestrial Ecosystem Model (TEM)

The TEM is a process-based ecosystem model that uses spatially referenced information on climate, elevation, soils, vegetation and water availability to estimate monthly vegetation and soil carbon and nitrogen fluxes and pool sizes. TEM is well-documented and has been used to examine patterns of terrestrial carbon dynamics across the globe including how they are influenced by multiple factors such as CO<sub>2</sub> fertilization, climate change and variability, land-use change, and ozone pollution (Melillo *et al.*, 1993; McGuire *et al.*, 1997, 2000a,b, 2001; Tian *et al.*, 1998, 1999, 2000, 2003; Xiao *et al.*, 1997, 1998; Prinn *et al.*, 1999; Reilly *et al.*, 1999, 2007a,b; Clein *et al.*, 2000; Webster *et al.*, 2003; Felzer *et al.*, 2004, 2005, 2007; Brovkin *et al.*, 2006; Sokolov *et al.*, 2008).

To determine the influence of land use change on terrestrial carbon dynamics, we calculate the net carbon exchange (NCE) between terrestrial ecosystems and the atmosphere by accounting for the carbon gained or lost due to ecosystem metabolism, as represented by net ecosystem production (NEP), the carbon lost during the conversion of natural ecosystems to agriculture ( $E_C$ ) and the carbon lost during the decomposition of agricultural and wood products ( $E_P$ ) as follows:

$$\text{NCE} = \text{NEP} - E_C - E_P \tag{A1}$$

Net ecosystem production is the balance between the uptake of carbon by vegetation to produce biomass and the release of carbon from respiration of living organisms and decomposition of dead organic matter within an ecosystem. A positive value of NCE represents carbon sequestration by terrestrial ecosystems whereas a negative value means that terrestrial

ecosystems are losing carbon. Further details of these TEM calculations may be found elsewhere (McGuire *et al.*, 2001; Tian *et al.*, 2003; Felzer *et al.*, 2004).

To simulate the carbon, nitrogen and water dynamics of cellulosic biofuels, we use the extant grassland parameterization of TEM to represent a generic cellulosic biofuel crop in a manner similar to that used by Felzer *et al.* (2004, 2005) for row-crop agriculture. In this study, we assume that both food crops and biofuel crops are optimally fertilized so that the productivity of these crops do not experience any nitrogen limitations.

Recently, a dynamic cohort approach has been adopted to represent the influence of land-use change on terrestrial carbon dynamics in TEM. In this approach, TEM initially assumes a  $0.5^\circ$  latitude x  $0.5^\circ$  longitude grid cell is covered by undisturbed potential vegetation, which is represented by an initial cohort that is assigned the entire land area of the grid cell. When a disturbance occurs, a new cohort is formed and a certain amount of land area within the grid cell is then subtracted from the undisturbed potential vegetation cohort and assigned to the new disturbed cohort. Disturbance-related carbon fluxes from the terrestrial ecosystem are calculated and the terrestrial carbon stocks are adjusted within the new disturbed cohort to account for the initial effect of the disturbance. The TEM is then used to simulate the recovery of terrestrial carbon dynamics after a disturbance within the context of local environmental conditions for the new disturbed cohort. As time progresses in the TEM simulation and more disturbances occur, more cohorts are added to the grid cell. As each disturbance and its effects are tracked separately within TEM, different types of disturbances within a grid cell can be considered simultaneously and allows TEM to consider the impacts of multiple disturbances on terrestrial carbon and nitrogen dynamics. The timing, location and affected area of a disturbance are prescribed by a spatially-explicit time-series land cover data set such as that described by Hurtt *et al.* (2006).

## **A.2 MIT Emissions Predictions and Policy Analysis (EPPA) Model**

The MIT Emissions Prediction and Policy Analysis (EPPA) model is a recursive-dynamic multi-regional computable general equilibrium (CGE) model of the world economy (Jacoby *et al.*, 1997; Babiker *et al.*, 2001; Paltsev *et al.*, 2005; Gurgel *et al.*, 2007; Reilly *et al.*, 2007a,b; Wang, 2008). The model is based on the Global Trade Analysis Project (GTAP) data base (Hertel, 1997; Dimaranan and McDougall, 2002) with the data aggregated into 16 regions and 24 sectors (**Table A1**). In the version of the model used here (EPPA4, Paltsev *et al.*, 2005, Gurgel *et*

*al.*, 2007; Wang, 2008), five of these sectors (**Table A2**) require land inputs that have been stratified into five land classes—cropland, pastureland, managed forest land, unmanaged grasslands, and unmanaged forest. Managed forests are those forests which have been disturbed by timber harvest since 1700. Unmanaged grasslands and forests do not produce any goods in the economy, but do provide recreational and conservation services (Wang, 2008). Conversion among these land classes is driven by economics, which consider the competition for land among alternative uses including the production of food, biofuels and wood products. The EPPA model also incorporates United States EPA inventory data and projections on greenhouse gas (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, HFCs, PFCs, and SF<sub>6</sub>) and air pollutant emissions (SO<sub>2</sub>, NO<sub>x</sub>, black carbon, organic carbon, NH<sub>3</sub>, CO, VOC) to estimate anthropogenic emissions of these compounds. These emission estimates may then be used to determine effects on atmospheric composition, climate and productivity of terrestrial ecosystems. The EPPA model projects the global economy, land use, and associated anthropogenic emissions into the future using a 5-year time step.

To enable linkages to TEM, each of the five land classes in each of the 16 EPPA regions has been assigned a unit price (i.e., the ratio of total land value over total land area) based on a comparison of the distribution of land cover in 1997 as described by the Hurtt *et al.* (2006) data set to the corresponding GTAP land-value data of cropland, pastured and managed forest (Lee *et al.*, 2005). For unmanaged forests, where services and prices are not explicitly reflected in normal economic accounting, we use data on access costs and timber output from the Global Timber Market and Forestry data Project (Sohngen, 2007) to assess the value of potential future harvest of the stock of standing timber and the residual value of land from future regrowth and harvest. The unit price of unmanaged forest is assumed to be the value of future harvests once the timber stock is gone, assuming that the value of the land rests in its ability to produce future harvests. The price ratio of unmanaged forest to managed forest is then applied to the price of pastures to obtain the unit price for unmanaged grasslands. The unit price of each land type is then used to determine changes in the land area required to support future market demand for food, biofuels and wood products based on associated changes in land value. The land use change is represented by economic relations that produce higher productivity land from lower productivity land by adding inputs—essentially investing in land improvements. Two versions of the model were developed—one that simply allowed any conversion that was economic, and a second that limited conversion through an elasticity of substitution that was based on observed

willingness to convert natural land to agricultural use. As the first version tended to lead to more conversion than we would expect based on historical evidence, we called it the “deforestation” scenario. The second approach was called the “intensification” scenario, since it is based on land supply elasticities calculated from agricultural land expansion observed in last decades. Further details of this approach are provided in Gurgel *et al.* (2007).

**Table A1.** Regions and Sectors in the EPPA4 Model. (Paltsev *et al.*, 2005).

<b>Country/Region</b>	<b>Sectors</b>
<b>Annex B</b>	<b>Non-Energy</b>
United States (USA)	Crops (CROP)
Canada (CAN)	Livestock (LIVE)
Japan (JPN)	Forestry (FORS)
European Union+ (EUR)	Food (FOOD)
Australia/New Zealand (ANZ)	Services (SERV)
Former Soviet Union (FSU)	Energy Intensive Products (EINT)
Eastern Europe (EET)	Other Industries Products (OTHR)
	Industrial Transportation (TRAN)
	Household Transportation (HTRN)
<b>Non-Annex B</b>	<b>Energy</b>
India (IND)	Coal (COAL)
China (CHN)	Crude Oil (OIL)
Indonesia (IDZ)	Refined Oil (ROIL)
Higher Income East Asia (ASI)	Natural Gas (GAS)
Mexico (MEX)	Electric: Fossil (ELEC)
Central and South America (LAM)	Electric: Hydro (HYDR)
Middle East (MES)	Electric: Nuclear (NUCL)
Africa (AFR)	Advanced Energy Technologies
Rest of World (ROW)	Electric: Biomass (BELE)
	Electric: Natural Gas Combined Cycle (NGCC)
	Electric: NGCC with CO <sub>2</sub> Capture and Storage (NGCAP)
	Electric: Integrated Coal Gasification with CO <sub>2</sub> Capture and Storage (IGCAP)
	Electric: Solar and Wind (SOLW)
	Liquid Fuel from Biomass (BOIL)
	Oil from Shale (SYNO)
	Synthetic Gas from Coal (SYNG)

**Table A2.** Sectors requiring land inputs (see also Wang, 2008).

<b>Sector</b>	<b>Land-use class</b>
Crops (CROP)	Cropland
Forestry (FORS)	Managed Forests
Livestock (LIVE)	Pasture
Electric: biomass (BELE)	Cropland
Liquid fuel from biomass (BOIL)	Cropland

A detailed description of the biofuels sector in EPPA can be found in Gurgel *et al.* (2007). The sector considers the growth and conversion of a cellulosic crop to a liquid fuel, which is a perfect substitute for refined oil. Maximum biomass production, in dry tons per hectare per year ( $\text{t ha}^{-1} \text{yr}^{-1}$ ), is assumed to vary by region, with highest initial productivity in Latin America ( $15 \text{ t ha}^{-1} \text{yr}^{-1}$ ), and lowest productivity in Canada ( $3 \text{ t ha}^{-1} \text{yr}^{-1}$ ). Other regions fall between these extremes, reflecting climatological limits to growth imposed by low moisture or temperature. Productivity of land in each region is allowed to change over time due to both technological improvement, and relative changes in NPP from the reference year, as simulated by TEM. Technological improvement include the possibility of both increased yield from biomass crops (1% increase per year) and increased efficiencies of cellulosic conversion processes. The energy embodied by biomass is considered to be 20 GJ per dry ton of biomass, with a conversion efficiency of 40% to liquid fuel. As a consequence, initial net energy yield per hectare ranges from  $120 \text{ GJ ha}^{-1} \text{yr}^{-1}$  in Latin America down to  $24 \text{ GJ ha}^{-1} \text{yr}^{-1}$  in Canada.

### **A.3 Downscaling EPPA Estimates of Land-use Change**

Since the EPPA model estimates land-use changes at the regional scale in response to economic forces, the regional changes estimated by EPPA need to be distributed to the  $0.5^\circ \times 0.5^\circ$  grid level before this information can be used by TEM. To downscale this information, assume that the spatial distribution of land-use change is determined by the spatial variations in land cover type, land productivity, climate, and management decisions across and within the EPPA regions. After assigning each grid cell to an EPPA region based on political boundaries (**Table A3**), we have developed a statistical approach based on an adaptation of the mixed-entropy method of You and Wood (2006) that takes into account the effects of climate conditions, farmer's experience of land productivities and human accessibility on land-use changes. As in their approach, we use some information to estimate a "prior" land-use share (i.e. proportion of a grid cell under a particular land use) for each grid cell. Later, we adjust this prior

to assure that the sum of areas in all grid cells within an EPPA region is consistent with regional EPPA shares. Basically, we seek to econometrically estimate the equation:

$$S_{l,n} = \alpha_l + \sum_i \beta_{l,i} NPP_{l,n} + \sum_i \chi_{l,i} NPP_{l,n}^2 + \delta_l T_n + \phi_l T_n^2 + \varphi_l P_n + \gamma_l P_n^2 + \eta_l D_n + \varepsilon_l, \quad (\text{A2})$$

where: subscripts  $l$  and  $i$  represent the land use categories (i.e., cropland, pasture, managed forest, unmanaged forest and unmanaged grassland),  $n$  represents the grid cell,  $S_{l,n}$  is the share of land use class  $l$  in grid cell  $n$ ,  $NPP_{l,n}$  is the averaged net primary productivity of the prior 5 years as estimated by TEM,  $T_n$  is the surface air temperature of grid cell  $n$ ,  $P_n$  is the precipitation in grid cell  $n$ ,  $D_n$  is distance between the center of grid cell  $n$  and the closest urban area. The parameters associated with each explanatory variable are represented by  $\beta_{l,i}$ ,  $\chi_{l,i}$ ,  $\delta_l$ ,  $\phi_l$ ,  $\varphi_l$ ,  $\gamma_l$  and  $\eta_l$ .  $\alpha_l$  is the linear intercept and  $\varepsilon_l$  is the error term. Equation A2 is estimated for each land-use class in each EPPA region, based on historical and simulated data for the period 1970 to 2000.

The historical data about  $S_{l,n}$  are obtained from Hurtt *et al.* (2006),  $NPP_{l,n}$  is determined by TEM simulations using the climate conditions and the Hurtt *et al.* (2006) land cover for this time period, historical grid data about  $T_n$  is from Brohan *et al.* (2006) and  $P_n$  from Hulme *et al.* (1998), and  $D_n$  data was calculated from Demographia (2007). Besides  $D_n$ , all other independent variables enter equation 1 as linear and squared terms (if significant). The NPP of all land classes affects the share of a particular land use class because the decision of changing a particular land-use has implications on other land classes in the grid cell. Human accessibility affects the decisions of land-use allocation because costs of transportation and access to inputs and markets decrease the likelihood a grid cell will be chosen for agriculture if it is far from an urban area. As emissions of greenhouse gases and other pollutants affect both climate and land productivity, equation A2 allows decisions of land use allocation to be influenced by climate policy.



**Table A3.** Association of EPPA4 regions to countries and territories across the globe.

<b>EPPA region</b>	<b>Countries and Territories</b>
AFR	Algeria, Angola, Benin, Botswana, Burkino Faso, Burundi, Cameroon, Canary Islands, Cape Verde, Central African Republic, Chad, Comoros, Democratic Republic of Congo, Djibouti, Egypt, Equatorial Guinea, Eritrea, Ethiopia, Europa Island, Gabon, Gambia, Ghana, Glorioso Islands, Guinea, Guinea-Bissau, Ivory Coast, Juan De Nova Island, Kenya, Lesotho, Liberia, Libya, Madagascar, Madeira, Malawi, Mali, Mauritania, Mauritius, Mayotte, Morocco, Mozambique, Namibia, Niger, Nigeria, Republic of Congo, Reunion, Rwanda, Saint Helena, Sao Tome and Principe, Senegal, Seychelles, Sierra Leone, Somalia, South Africa, Sudan, Swaziland, Tanzania, Togo, Tromelin Island, Tunisia, Uganda, Western Sahara, Zambia, Zimbabwe
ANZ	Australia, Cook Islands, New Zealand, Niue, Norfolk Island, Tokelau
ASI	Malaysia, Philippines, Singapore, South Korea, Taiwan, Thailand
CAN	Canada
CHN	China, Hong Kong, Paracel Islands
EET	Bulgaria, Czech Republic, Hungary, Poland, Romania, Slovakia, Slovenia
EUR	Austria, Belgium, Denmark, Faroe Islands, Finland, France, Germany, Greece, Iceland, Ireland, Italy, Liechtenstein, Luxembourg, Malta, The Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom
FSU	Armenia, Azerbaijan, Belarus, Estonia, Georgia, Kazakhstan, Kyrgyzstan, Latvia, Lithuania, Moldova, Russia, Tajikistan, Turkmenistan, Ukraine, Uzbekistan
IDZ	Indonesia, Timor Leste
IND	India
JPN	Japan
LAM	Anguilla, Antigua and Barbuda, Argentina, Aruba, Bahamas, Barbados, Belize, Bermuda, Bolivia, Brazil, Cayman Islands, Chile, Colombia, Costa Rica, Cuba, Dominica, Dominican Republic, Ecuador, El Salvador, Falkland Islands, French Guiana, Grenada, Guadeloupe, Guatemala, Guyana, Haiti, Honduras, Jamaica, Martinique, Montserrat, Netherland Antilles, Nicaragua, Panama, Paraguay, Peru, Puerto Rico, Saint Kitts and Nevis, Saint Lucia, Saint Pierre and Miquelon, Saint Vincent and the Grenadines, Suriname, Trinidad and Tobago, Turks and Caicos Islands, Uruguay, Venezuela, Virgin Islands
MES	Bahrain, Iran, Iraq, Israel, Jordan, Kuwait, Lebanon, Oman, Palestinian Territories, Qatar, Saudi Arabia, Syria, United Arab Emirates, Yemen
MEX	Mexico
ROW	Afghanistan, Albania, American Samoa, Bangladesh, Bhutan, Bosnia-Herzegovina, British Indian Ocean Territory, Brunei, Cambodia, Croatia, Cyprus, Fiji, French Polynesia, French Southern and Antarctic Lands, Futuna Island, Greenland, Guam, Kiribati, Laos, Macedonia, Maldives, Marshall Islands, Micronesia, Mongolia, Montenegro, Myanmar, Nauru, Nepal, New Caledonia, Northern Mariana Islands, North Korea, Pakistan, Palau, Papua New Guinea, Pitcairn Islands, Samoa, Serbia, Solomon Islands, Sri Lanka, South Georgia Island, Tonga, Turkey, Tuvalu, Vanuatu, Vietnam, Wallis Island
USA	United States of America

The historical cross-section land-use allocation decisions captured by the estimation of the parameters for equation A2 are then used to forecast the priors of gridded land use in the future using estimates of NPP projected by TEM and estimates of surface air temperature and precipitation projected by the MIT IGSM downscaled to grid level. Urban areas are assumed to remain constant over this period. Because cellulosic biofuels production is not present in the historical data, we determined a prior for biofuels by setting the share of land covered by cellulosic biofuels as the regional share of biofuels forecasted by EPPA in all grid cells where cropland area is at least 5%. This means that cellulosic biofuels are distributed equally across cropland areas in a region.

The land-use priors projected by the downscaling statistical model are then compared to the regional land shares from EPPA and consistently corrected to match them. This correction is done to assure that the sum of shares in each grid cell equals to one and, in each region, the sum of the areas of a particular land use class over all grid cells match the area predicted by EPPA. Both consistency conditions are assured simply re-scaling grid cells until these conditions are met. After downscaling, changes in land shares estimated by EPPA need to be mapped to the disturbance cohorts used by TEM.

#### **A.4 Mapping EPPA Land Shares to TEM Cohorts**

To represent contemporary land cover in this study, TEM uses the IGSMVEG classification described by Schlosser *et al.* (2007) and information from Hurtt *et al.* (2006) that describes annual changes in land cover from 1700 to 2000. The IGSMVEG classification stratifies global vegetation into 35 upland and wetland cover types and a 0.5° latitude x 0.5° longitude grid cell may be covered by a mosaic of land cover types based on spatially-explicit information from Melillo *et al.* (1993), Bonan *et al.* (2002) and Matthews and Fung (1987). The Hurtt *et al.* (2006) data describes spatially-explicit transitions among primary vegetation, secondary vegetation, croplands and pastures. Primary vegetation is land cover that has not been directly disturbed by human activities. Secondary vegetation is a result of disturbance to primary vegetation (e.g., timber harvest or wood gathering) or abandonment of croplands or pastures. As described earlier, TEM assumes a grid cell is initially represented by a set of cohorts that describe the distribution of “potential” vegetation land cover before any disturbance occurs. The spatially-explicit transition data from Hurtt *et al.* (2006) is then used to determine the timing and location of the creation of disturbance cohorts and the introduction of new food crop, biofuel crop, and pasture

land cover types to each grid cell. The data set also determines the area of the grid cell that is affected by these transitions and this area is subtracted from the area of the appropriate existing cohorts. As a result of this approach of tracking land-use history, a 0.5° latitude x 0.5° longitude grid cell may have up to 1210 cohorts by the year 2000. The distribution of TEM cohorts during 2000, is then used as the initial land cover for developing future land use scenarios using changes in EPPA land shares.

To determine future transitions in land cover, differences in the downscaled land shares estimated by EPPA are determined for successive 5-year time steps, interpolated to annual time steps and then mapped to the TEM cohort structure. The resulting spatially-explicit time-series transition data set includes transitions among croplands used to grow food crops, croplands used to grow cellulosic biofuels, pastures, managed forests, unmanaged forests, and unmanaged grasslands. The EPPA-derived transitions also indicate the timing and location of timber harvests in the managed forests.

For timber harvests, we assume that a fraction (i.e.,  $1/rotationage$ ) of managed forests in a grid cell will be harvested each year based on the rotation age determined for those forests. Rotation age is assumed to vary with latitude as follows:

$$rotationage = 9 + 0.5|latitude|^{1.1} \quad (A3)$$

Thus, short rotation ages (e.g., 9 years) are assumed to occur near the equator and longer rotation ages (e.g., 60 years near the Arctic Circle) are assumed to occur with distance away from the equator in general agreement with other studies (**Table A4**).

The EPPA-derived transitions are mapped to the IGSMVEG types used by TEM (**Table A5**) based on the vegetation cover within the grid cell and used to create new disturbance cohorts into the future. No changes are assumed to occur in the distribution of tundra, wetlands, salt marshes, or deserts. The new spatially-explicit time-series cohort data set is then used to prescribe annual changes in land cover for TEM projections of future terrestrial carbon fluxes.

### **A.5 Development of the Climate Scenario**

As biofuels are being promoted as an important part of the global energy mix to meet the climate change challenge, we have developed a single climate scenario based on a particular climate policy to examine the potential effects of alternative aggressive future cellulosic biofuels

programs on the terrestrial biosphere within the context of concurrent climate change. The climate scenario is based on a policy to control greenhouse gas emissions from industrial and fossil fuel sources that would stabilize the atmosphere's CO<sub>2</sub> concentration at 550 ppmv (Paltsev *et al.*, 2008). Under this climate policy, developed countries would gradually phase in a 50% reduction in emissions by 2050, like that suggested in recent G8 meetings and consistent with proposed goals in Europe and in pending bills before the U.S. Congress. Developing countries would delay their mitigation action until 2025, and intensify reductions in 2035. Similar to the provisions of other existing climate policies, fossil fuel emissions of CO<sub>2</sub>, including those resulting from production of biofuels, are assumed to be controlled, but emissions from land-use change are not. As a result, the climate policy scenario used here does not provide incentives to avoid land-use emissions resulting from land clearing to produce biofuels.

**Table A4.** Examples of rotation age used for timber harvest by this study in comparison to the optimal rotation age from Sohngen *et al.* (1999).


<b>Latitude</b>	<b>This Study's Rotation Age (years)</b>	<b>Sohngen's Rotation Age (years)</b>
15° N or S	19	20 for southern pine plantations 9 for southern eucalyptus plantations
30° N or S	30	31 to 40 for natural and plantation pine plantations in the southern USA 12 for eucalyptus 30 for pine plantations in Oceania 27 for plantations in southern China 11 for plantations in the Iberian peninsula 17 for pine plantations in southern Central Asia
45° N or S	42	49 to 60 for softwood forests in Eastern provinces and softwood and hardwood forests in Lake Provinces of Canada 30 for temperate forests in Central Asia
60° N or S	54	54 for Nordic plantations in Europe 45 to 70 for US Pacific Northwest Douglas fir-hemlock forests, Western Pine, North-Eastern and Great Lakes softwood general type, oak-hickory and maple-beech-birch forests. 84 to 91 for conifers, temperate hardwoods and boreal hardwoods in Russia

**Table A5.** Relationship of EPPA land shares to IGSMVEG vegetation types (Schlosser *et al.*, 2007)


<b>EPPA Sector/land-use class</b>	<b>IGSMVEG</b>
CROP Cropland	Crop 1
BOIL Cropland	Crop 2
Pasture	Pasture
Managed Forests, Unmanaged forests	Needle-leaf Evergreen Tree (NET) temperate, Needle-leaf Evergreen Tree (NET) boreal, Needle-leaf Deciduous Tree (NDT) boreal, Broadleaved Evergreen Tree (BET) tropical, Broadleaved Evergreen Tree (BET) temperate, Broadleaved Deciduous Tree (BDT) tropical, Broadleaved Deciduous Tree (BDT) temperate, Broadleaved Deciduous Tree (BDT) boreal, Broadleaved Evergreen Shrub (BES) temperate, Broadleaved Deciduous Shrub (BDS) temperate, Broadleaved Deciduous Shrub (BDS) boreal, Floodplains (Tree tropical), Floodplains (Tree temperate)
Unmanaged grasslands	C3 grass, C4 grass, Floodplains (No-tree tropical), Floodplains (No-tree temperate)

The GHG and other pollutant emissions projected by EPPA based on this climate policy have been used to drive the coupled atmospheric and climate module within the MIT IGSM to estimate zonal (i.e., 4° latitudinal bands) changes in atmospheric composition and climate over the 21<sup>st</sup> century. In the resulting climate scenario, atmospheric CO<sub>2</sub> concentrations increase by 163 ppmv; the global mean AOT40 ozone index, a measure of the accumulated hourly ozone levels above a threshold of 40 ppb, almost doubles; global mean air temperatures increases by 2.4°; and global mean precipitation increases by 30 mm yr<sup>-1</sup> by 2100 (**Figure A1**). The monthly zonal changes in climate are distributed to the 0.5° x 0.5° spatial resolution by applying these changes to a baseline climate (Cramer and Leemans, 2001) as described previously by Xiao *et al.* (1997). Zonal changes in the AOT40 index have been downscaled as described by Felzer *et al.* (2005). The downscaled climate is then used to downscale the land-use changes projected by EPPA as described in section A.3 and to drive TEM to develop gridded estimates of net carbon exchange between terrestrial ecosystems and the atmosphere.


a) Atmospheric CO<sub>2</sub> concentrations




b) AOT40 ozone index



c) Global mean air temperature



d) Global mean precipitation



**Figure A1.** Projected changes in environmental factors from the climate policy including: a) global atmospheric CO<sub>2</sub> concentrations, b) mean global AOT40 ozone index, c) global mean annual air temperatures, and d) global annual precipitation.

## REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

1. **Uncertainty in Climate Change Policy Analysis**  
*Jacoby & Prinn* December 1994
2. **Description and Validation of the MIT Version of the GISS 2D Model** *Sokolov & Stone* June 1995
3. **Responses of Primary Production and Carbon Storage to Changes in Climate and Atmospheric CO<sub>2</sub> Concentration** *Xiao et al.* October 1995
4. **Application of the Probabilistic Collocation Method for an Uncertainty Analysis** *Webster et al.* January 1996
5. **World Energy Consumption and CO<sub>2</sub> Emissions: 1950-2050** *Schmalensee et al.* April 1996
6. **The MIT Emission Prediction and Policy Analysis (EPPA) Model** *Yang et al.* May 1996 (*superseded* by No. 125)
7. **Integrated Global System Model for Climate Policy Analysis** *Prinn et al.* June 1996 (*superseded* by No. 124)
8. **Relative Roles of Changes in CO<sub>2</sub> and Climate to Equilibrium Responses of Net Primary Production and Carbon Storage** *Xiao et al.* June 1996
9. **CO<sub>2</sub> Emissions Limits: Economic Adjustments and the Distribution of Burdens** *Jacoby et al.* July 1997
10. **Modeling the Emissions of N<sub>2</sub>O and CH<sub>4</sub> from the Terrestrial Biosphere to the Atmosphere** *Liu* Aug. 1996
11. **Global Warming Projections: Sensitivity to Deep Ocean Mixing** *Sokolov & Stone* September 1996
12. **Net Primary Production of Ecosystems in China and its Equilibrium Responses to Climate Changes** *Xiao et al.* November 1996
13. **Greenhouse Policy Architectures and Institutions** *Schmalensee* November 1996
14. **What Does Stabilizing Greenhouse Gas Concentrations Mean?** *Jacoby et al.* November 1996
15. **Economic Assessment of CO<sub>2</sub> Capture and Disposal** *Eckaus et al.* December 1996
16. **What Drives Deforestation in the Brazilian Amazon?** *Pfaff* December 1996
17. **A Flexible Climate Model For Use In Integrated Assessments** *Sokolov & Stone* March 1997
18. **Transient Climate Change and Potential Croplands of the World in the 21st Century** *Xiao et al.* May 1997
19. **Joint Implementation: Lessons from Title IV's Voluntary Compliance Programs** *Atkeson* June 1997
20. **Parameterization of Urban Subgrid Scale Processes in Global Atm. Chemistry Models** *Calbo et al.* July 1997
21. **Needed: A Realistic Strategy for Global Warming** *Jacoby, Prinn & Schmalensee* August 1997
22. **Same Science, Differing Policies; The Saga of Global Climate Change** *Skolnikoff* August 1997
23. **Uncertainty in the Oceanic Heat and Carbon Uptake and their Impact on Climate Projections** *Sokolov et al.* September 1997
24. **A Global Interactive Chemistry and Climate Model** *Wang, Prinn & Sokolov* September 1997
25. **Interactions Among Emissions, Atmospheric Chemistry & Climate Change** *Wang & Prinn* Sept. 1997
26. **Necessary Conditions for Stabilization Agreements** *Yang & Jacoby* October 1997
27. **Annex I Differentiation Proposals: Implications for Welfare, Equity and Policy** *Reiner & Jacoby* Oct. 1997
28. **Transient Climate Change and Net Ecosystem Production of the Terrestrial Biosphere** *Xiao et al.* November 1997
29. **Analysis of CO<sub>2</sub> Emissions from Fossil Fuel in Korea: 1961-1994** *Choi* November 1997
30. **Uncertainty in Future Carbon Emissions: A Preliminary Exploration** *Webster* November 1997
31. **Beyond Emissions Paths: Rethinking the Climate Impacts of Emissions Protocols** *Webster & Reiner* November 1997
32. **Kyoto's Unfinished Business** *Jacoby et al.* June 1998
33. **Economic Development and the Structure of the Demand for Commercial Energy** *Judson et al.* April 1998
34. **Combined Effects of Anthropogenic Emissions and Resultant Climatic Changes on Atmospheric OH** *Wang & Prinn* April 1998
35. **Impact of Emissions, Chemistry, and Climate on Atmospheric Carbon Monoxide** *Wang & Prinn* April 1998
36. **Integrated Global System Model for Climate Policy Assessment: Feedbacks and Sensitivity Studies** *Prinn et al.* June 1998
37. **Quantifying the Uncertainty in Climate Predictions** *Webster & Sokolov* July 1998
38. **Sequential Climate Decisions Under Uncertainty: An Integrated Framework** *Valverde et al.* September 1998
39. **Uncertainty in Atmospheric CO<sub>2</sub> (Ocean Carbon Cycle Model Analysis)** *Holian* Oct. 1998 (*superseded* by No. 80)
40. **Analysis of Post-Kyoto CO<sub>2</sub> Emissions Trading Using Marginal Abatement Curves** *Ellerman & Decaux* Oct. 1998
41. **The Effects on Developing Countries of the Kyoto Protocol and CO<sub>2</sub> Emissions Trading** *Ellerman et al.* November 1998
42. **Obstacles to Global CO<sub>2</sub> Trading: A Familiar Problem** *Ellerman* November 1998
43. **The Uses and Misuses of Technology Development as a Component of Climate Policy** *Jacoby* November 1998
44. **Primary Aluminum Production: Climate Policy, Emissions and Costs** *Harnisch et al.* December 1998
45. **Multi-Gas Assessment of the Kyoto Protocol** *Reilly et al.* January 1999
46. **From Science to Policy: The Science-Related Politics of Climate Change Policy in the U.S.** *Skolnikoff* January 1999
47. **Constraining Uncertainties in Climate Models Using Climate Change Detection Techniques** *Forest et al.* April 1999
48. **Adjusting to Policy Expectations in Climate Change Modeling** *Shackley et al.* May 1999
49. **Toward a Useful Architecture for Climate Change Negotiations** *Jacoby et al.* May 1999
50. **A Study of the Effects of Natural Fertility, Weather and Productive Inputs in Chinese Agriculture** *Eckaus & Tso* July 1999
51. **Japanese Nuclear Power and the Kyoto Agreement** *Babiker, Reilly & Ellerman* August 1999
52. **Interactive Chemistry and Climate Models in Global Change Studies** *Wang & Prinn* September 1999
53. **Developing Country Effects of Kyoto-Type Emissions Restrictions** *Babiker & Jacoby* October 1999

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

## REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

54. **Model Estimates of the Mass Balance of the Greenland and Antarctic Ice Sheets** *Bugnion* Oct 1999
55. **Changes in Sea-Level Associated with Modifications of Ice Sheets over 21st Century** *Bugnion* October 1999
56. **The Kyoto Protocol and Developing Countries** *Babiker et al.* October 1999
57. **Can EPA Regulate Greenhouse Gases Before the Senate Ratifies the Kyoto Protocol?** *Bugnion & Reiner* November 1999
58. **Multiple Gas Control Under the Kyoto Agreement** *Reilly, Mayer & Harnisch* March 2000
59. **Supplementarity: An Invitation for Monopsony?** *Ellerman & Sue Wing* April 2000
60. **A Coupled Atmosphere-Ocean Model of Intermediate Complexity** *Kamenkovich et al.* May 2000
61. **Effects of Differentiating Climate Policy by Sector: A U.S. Example** *Babiker et al.* May 2000
62. **Constraining Climate Model Properties Using Optimal Fingerprint Detection Methods** *Forest et al.* May 2000
63. **Linking Local Air Pollution to Global Chemistry and Climate** *Mayer et al.* June 2000
64. **The Effects of Changing Consumption Patterns on the Costs of Emission Restrictions** *Lahiri et al.* Aug 2000
65. **Rethinking the Kyoto Emissions Targets** *Babiker & Eckaus* August 2000
66. **Fair Trade and Harmonization of Climate Change Policies in Europe** *Viguié* September 2000
67. **The Curious Role of "Learning" in Climate Policy: Should We Wait for More Data?** *Webster* October 2000
68. **How to Think About Human Influence on Climate** *Forest, Stone & Jacoby* October 2000
69. **Tradable Permits for Greenhouse Gas Emissions: A primer with reference to Europe** *Ellerman* Nov 2000
70. **Carbon Emissions and The Kyoto Commitment in the European Union** *Viguié et al.* February 2001
71. **The MIT Emissions Prediction and Policy Analysis Model: Revisions, Sensitivities and Results** *Babiker et al.* February 2001 (*superseded* by No. 125)
72. **Cap and Trade Policies in the Presence of Monopoly and Distortionary Taxation** *Fullerton & Metcalf* March '01
73. **Uncertainty Analysis of Global Climate Change Projections** *Webster et al.* Mar. '01 (*superseded* by No. 95)
74. **The Welfare Costs of Hybrid Carbon Policies in the European Union** *Babiker et al.* June 2001
75. **Feedbacks Affecting the Response of the Thermohaline Circulation to Increasing CO<sub>2</sub>** *Kamenkovich et al.* July 2001
76. **CO<sub>2</sub> Abatement by Multi-fueled Electric Utilities: An Analysis Based on Japanese Data** *Ellerman & Tsukada* July 2001
77. **Comparing Greenhouse Gases** *Reilly et al.* July 2001
78. **Quantifying Uncertainties in Climate System Properties using Recent Climate Observations** *Forest et al.* July 2001
79. **Uncertainty in Emissions Projections for Climate Models** *Webster et al.* August 2001
80. **Uncertainty in Atmospheric CO<sub>2</sub> Predictions from a Global Ocean Carbon Cycle Model** *Holian et al.* September 2001
81. **A Comparison of the Behavior of AO GCMs in Transient Climate Change Experiments** *Sokolov et al.* December 2001
82. **The Evolution of a Climate Regime: Kyoto to Marrakech** *Babiker, Jacoby & Reiner* February 2002
83. **The "Safety Valve" and Climate Policy** *Jacoby & Ellerman* February 2002
84. **A Modeling Study on the Climate Impacts of Black Carbon Aerosols** *Wang* March 2002
85. **Tax Distortions and Global Climate Policy** *Babiker et al.* May 2002
86. **Incentive-based Approaches for Mitigating Greenhouse Gas Emissions: Issues and Prospects for India** *Gupta* June 2002
87. **Deep-Ocean Heat Uptake in an Ocean GCM with Idealized Geometry** *Huang, Stone & Hill* September 2002
88. **The Deep-Ocean Heat Uptake in Transient Climate Change** *Huang et al.* September 2002
89. **Representing Energy Technologies in Top-down Economic Models using Bottom-up Information** *McFarland et al.* October 2002
90. **Ozone Effects on Net Primary Production and Carbon Sequestration in the U.S. Using a Biogeochemistry Model** *Felzer et al.* November 2002
91. **Exclusionary Manipulation of Carbon Permit Markets: A Laboratory Test** *Carlén* November 2002
92. **An Issue of Permanence: Assessing the Effectiveness of Temporary Carbon Storage** *Herzog et al.* December 2002
93. **Is International Emissions Trading Always Beneficial?** *Babiker et al.* December 2002
94. **Modeling Non-CO<sub>2</sub> Greenhouse Gas Abatement** *Hyman et al.* December 2002
95. **Uncertainty Analysis of Climate Change and Policy Response** *Webster et al.* December 2002
96. **Market Power in International Carbon Emissions Trading: A Laboratory Test** *Carlén* January 2003
97. **Emissions Trading to Reduce Greenhouse Gas Emissions in the United States: The McCain-Lieberman Proposal** *Paltsev et al.* June 2003
98. **Russia's Role in the Kyoto Protocol** *Bernard et al.* Jun '03
99. **Thermohaline Circulation Stability: A Box Model Study** *Lucarini & Stone* June 2003
100. **Absolute vs. Intensity-Based Emissions Caps** *Ellerman & Sue Wing* July 2003
101. **Technology Detail in a Multi-Sector CGE Model: Transport Under Climate Policy** *Schafer & Jacoby* July 2003
102. **Induced Technical Change and the Cost of Climate Policy** *Sue Wing* September 2003
103. **Past and Future Effects of Ozone on Net Primary Production and Carbon Sequestration Using a Global Biogeochemical Model** *Felzer et al.* (revised) January 2004
104. **A Modeling Analysis of Methane Exchanges Between Alaskan Ecosystems and the Atmosphere** *Zhuang et al.* November 2003

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.



## REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

105. **Analysis of Strategies of Companies under Carbon Constraint** *Hashimoto* January 2004
106. **Climate Prediction: The Limits of Ocean Models** *Stone* February 2004
107. **Informing Climate Policy Given Incommensurable Benefits Estimates** *Jacoby* February 2004
108. **Methane Fluxes Between Terrestrial Ecosystems and the Atmosphere at High Latitudes During the Past Century** *Zhuang et al.* March 2004
109. **Sensitivity of Climate to Diapycnal Diffusivity in the Ocean** *Dalan et al.* May 2004
110. **Stabilization and Global Climate Policy** *Sarofim et al.* July 2004
111. **Technology and Technical Change in the MIT EPPA Model** *Jacoby et al.* July 2004
112. **The Cost of Kyoto Protocol Targets: The Case of Japan** *Paltsev et al.* July 2004
113. **Economic Benefits of Air Pollution Regulation in the USA: An Integrated Approach** *Yang et al.* (revised) Jan. 2005
114. **The Role of Non-CO<sub>2</sub> Greenhouse Gases in Climate Policy: Analysis Using the MIT IGSM** *Reilly et al.* Aug. '04
115. **Future U.S. Energy Security Concerns** *Deutch* Sep. '04
116. **Explaining Long-Run Changes in the Energy Intensity of the U.S. Economy** *Sue Wing* Sept. 2004
117. **Modeling the Transport Sector: The Role of Existing Fuel Taxes in Climate Policy** *Paltsev et al.* November 2004
118. **Effects of Air Pollution Control on Climate** *Prinn et al.* January 2005
119. **Does Model Sensitivity to Changes in CO<sub>2</sub> Provide a Measure of Sensitivity to the Forcing of Different Nature?** *Sokolov* March 2005
120. **What Should the Government Do To Encourage Technical Change in the Energy Sector?** *Deutch* May '05
121. **Climate Change Taxes and Energy Efficiency in Japan** *Kasahara et al.* May 2005
122. **A 3D Ocean-Seaice-Carbon Cycle Model and its Coupling to a 2D Atmospheric Model: Uses in Climate Change Studies** *Dutkiewicz et al.* (revised) November 2005
123. **Simulating the Spatial Distribution of Population and Emissions to 2100** *Asadoorian* May 2005
124. **MIT Integrated Global System Model (IGSM) Version 2: Model Description and Baseline Evaluation** *Sokolov et al.* July 2005
125. **The MIT Emissions Prediction and Policy Analysis (EPPA) Model: Version 4** *Paltsev et al.* August 2005
126. **Estimated PDFs of Climate System Properties Including Natural and Anthropogenic Forcings** *Forest et al.* September 2005
127. **An Analysis of the European Emission Trading Scheme** *Reilly & Paltsev* October 2005
128. **Evaluating the Use of Ocean Models of Different Complexity in Climate Change Studies** *Sokolov et al.* November 2005
129. **Future Carbon Regulations and Current Investments in Alternative Coal-Fired Power Plant Designs** *Sekar et al.* December 2005
130. **Absolute vs. Intensity Limits for CO<sub>2</sub> Emission Control: Performance Under Uncertainty** *Sue Wing et al.* January 2006
131. **The Economic Impacts of Climate Change: Evidence from Agricultural Profits and Random Fluctuations in Weather** *Deschenes & Greenstone* January 2006
132. **The Value of Emissions Trading** *Webster et al.* Feb. 2006
133. **Estimating Probability Distributions from Complex Models with Bifurcations: The Case of Ocean Circulation Collapse** *Webster et al.* March 2006
134. **Directed Technical Change and Climate Policy** *Otto et al.* April 2006
135. **Modeling Climate Feedbacks to Energy Demand: The Case of China** *Asadoorian et al.* June 2006
136. **Bringing Transportation into a Cap-and-Trade Regime** *Ellerman, Jacoby & Zimmerman* June 2006
137. **Unemployment Effects of Climate Policy** *Babiker & Eckaus* July 2006
138. **Energy Conservation in the United States: Understanding its Role in Climate Policy** *Metcalfe* Aug. '06
139. **Directed Technical Change and the Adoption of CO<sub>2</sub> Abatement Technology: The Case of CO<sub>2</sub> Capture and Storage** *Otto & Reilly* August 2006
140. **The Allocation of European Union Allowances: Lessons, Unifying Themes and General Principles** *Buchner et al.* October 2006
141. **Over-Allocation or Abatement? A preliminary analysis of the EU ETS based on the 2006 emissions data** *Ellerman & Buchner* December 2006
142. **Federal Tax Policy Towards Energy** *Metcalfe* Jan. 2007
143. **Technical Change, Investment and Energy Intensity** *Kratena* March 2007
144. **Heavier Crude, Changing Demand for Petroleum Fuels, Regional Climate Policy, and the Location of Upgrading Capacity** *Reilly et al.* April 2007
145. **Biomass Energy and Competition for Land** *Reilly & Paltsev* April 2007
146. **Assessment of U.S. Cap-and-Trade Proposals** *Paltsev et al.* April 2007
147. **A Global Land System Framework for Integrated Climate-Change Assessments** *Schlosser et al.* May 2007
148. **Relative Roles of Climate Sensitivity and Forcing in Defining the Ocean Circulation Response to Climate Change** *Scott et al.* May 2007
149. **Global Economic Effects of Changes in Crops, Pasture, and Forests due to Changing Climate, CO<sub>2</sub> and Ozone** *Reilly et al.* May 2007
150. **U.S. GHG Cap-and-Trade Proposals: Application of a Forward-Looking Computable General Equilibrium Model** *Gurgel et al.* June 2007
151. **Consequences of Considering Carbon/Nitrogen Interactions on the Feedbacks between Climate and the Terrestrial Carbon Cycle** *Sokolov et al.* June 2007
152. **Energy Scenarios for East Asia: 2005-2025** *Paltsev & Reilly* July 2007
153. **Climate Change, Mortality, and Adaptation: Evidence from Annual Fluctuations in Weather in the U.S.** *Deschênes & Greenstone* August 2007

## REPORT SERIES of the MIT *Joint Program on the Science and Policy of Global Change*

- 154. Modeling the Prospects for Hydrogen Powered Transportation Through 2100** *Sandoval et al.*  
February 2008
- 155. Potential Land Use Implications of a Global Biofuels Industry** *Gurgel et al.* March 2008
- 156. Estimating the Economic Cost of Sea-Level Rise**  
*Sugiyama et al.* April 2008
- 157. Constraining Climate Model Parameters from Observed 20<sup>th</sup> Century Changes** *Forest et al.* April 2008
- 158. Analysis of the Coal Sector under Carbon Constraints** *McFarland et al.* April 2008
- 159. Impact of Sulfur and Carbonaceous Emissions from International Shipping on Aerosol Distributions and Direct Radiative Forcing** *Wang & Kim* April 2008
- 160. Analysis of U.S. Greenhouse Gas Tax Proposals**  
*Metcalf et al.* April 2008
- 161. A Forward Looking Version of the MIT Emissions Prediction and Policy Analysis (EPPA) Model**  
*Babiker et al.* May 2008
- 162. The European Carbon Market in Action: Lessons from the first trading period** Interim Report  
*Convery, Ellerman, & de Perthuis* June 2008
- 163. The Influence on Climate Change of Differing Scenarios for Future Development Analyzed Using the MIT Integrated Global System Model** *Prinn et al.*  
September 2008
- 164. Marginal Abatement Costs and Marginal Welfare Costs for Greenhouse Gas Emissions Reductions: Results from the EPPA Model** *Holak et al.* November 2008
- 165. Uncertainty in Greenhouse Emissions and Costs of Atmospheric Stabilization** *Webster et al.* November 2008
- 166. Sensitivity of Climate Change Projections to Uncertainties in the Estimates of Observed Changes in Deep-Ocean Heat Content** *Sokolov et al.* November 2008
- 167. Sharing the Burden of GHG Reductions** *Jacoby et al.*  
November 2008
- 168. Unintended Environmental Consequences of a Global Biofuels Program** *Melillo et al.* January 2009